CAGTTTCTTCATCTGTAACATCAAATGAATAATAATACCAATCTCCTAGACTTCATAAGA GGATTAACAAAGACAAAATATGGGAAAAACATAACATGGCGTCCCATAATTATTAGATCT TATTATTGACACTAAAATGGCATTAAAATTACCAAAAGGAAGACAGCATCTGTTTCCTCT TTGGTCCTGAGCTGGTTAAAAGGAACACTGGTTGCCTGAACAGTCACACTTGCAACCATG **ATG**CCTAAACATTGCTTTCTAGGCTTCCTCATCAGTTTCTTCCTTACTGGTGTAGCAGGA ACTCAGTCAACGCATGAGTCTCTGAAGCCTCAGAGGGTACAATTTCAGTCCCGAAATTTT CACAACATTTTGCAATGGCAGCCTGGGAGGGCACTTACTGGCAACAGCAGTGTCTATTTT GTGCAGTACAAAATCATGTTCTCATGCAGCATGAAAAGCTCTCACCAGAAGCCAAGTGGA TGCTGGCAGCACATTTCTTGTAACTTCCCAGGCTGCAGAACATTGGCTAAATATGGACAG AGACAATGGAAAAATAAAGAAGACTGTTGGGGTACTCAAGAACTCTCTTGTGACCTTACC AGCTACTCAGAATGGAGCATGACGCCGCGGTTCACTCCCTGGTGGGAAACAAAAATAGAT CCTCCAGTCATGAATATAACCCAAGTCAATGGCTCTTTGTTGGTAATTCTCCATGCTCCA AATTTACCATATAGATACCAAAAGGAAAAAATGTATCTATAGAAGATTACTATGAACTA CTATACCGAGTTTTTATAATTAACAATTCACTAGAAAAGGAGCAAAAGGTTTATGAAGGG GCTCACAGAGCGGTTGAAATTGAAGCTCTAACACCACACTCCAGCTACTGTGTAGTGGCT GAAATATATCAGCCCATGTTAGACAGAAGAAGTCAGAGAAGTGAAGAGAGATGTGTGGAA ATTCCA**TGA**CTTGTGGAATTTGGCATTCAGCAATGTGGAAATTCTAAAGCTCCCTGAGAA CAGGATGACTCGTGTTTGAAGGATCTTATTTAAAATTGTTTTTTGTATTTTCTTAAAGCAA TATTCACTGTTACACCTTGGGGACTTCTTTGTTTATCCATTCTTTTATCCTTTATATTTC ATTTGTAAACTATATTTGAACGACATTCCCCCCGAAAAATTGAAATGTAAAGATGAGGCA

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA145887

><subunit 1 of 1, 262 aa, 1 stop

><MW: 30419, pI: 8.44, NX(S/T): 5

MPKHCFLGFLISFFLTGVAGTQSTHESLKPQRVQFQSRNFHNILQWQPGRALTGNSSVYF VQYKIMFSCSMKSSHQKPSGCWQHISCNFPGCRTLAKYGQRQWKNKEDCWGTQELSCDLT SETSDIQEPYYGRVRAASAGSYSEWSMTPRFTPWWETKIDPPVMNITQVNGSLLVILHAP NLPYRYQKEKNVSIEDYYELLYRVFIINNSLEKEQKVYEGAHRAVEIEALTPHSSYCVVA EIYQPMLDRRSQRSEERCVEIP

Important features of the protein:

Signal peptide:

Amino Acids

1-20

N-glycosylation sites:

Amino acids

55-59;165-169;170-174;191-195;208-212

N-myristoylation sites:

Amino acids

17-23;20-26;220-226

CTCAAGCATCACTTACAGGACCAGAGGGACAAGACATGACTGTGATGAGGAGCTGCTTTCGC CAATTTAACACCAAGAAGAATTGAGGCTGCTTGGGAGGAAGGCCAGGAGGAACACGAGACTG $\texttt{AGAG} \underline{\textbf{ATG}} \texttt{AATTTTCAACAGAGGCTGCAAAGCCTGTGGACTTTAGCCAGACCCTTCTGCCCTC}$ CTTTGCTGGCGACAGCCTCTCAAATGCAGATGGTTGTGCTCCCTTGCCTGGGTTTTACCCTG CTTCTCTGGAGCCAGGTATCAGGGGCCCAGGGCCAAGAATTCCACTTTGGGCCCTGCCAAGT GAAGGGGGTTGTTCCCCAGAAACTGTGGGAAGCCTTCTGGGCTGTGAAAGACACTATGCAAG CTCAGGATAACATCACGAGTGCCCGGCTGCTGCAGCAGGAGGTTCTGCAGAACGTCTCGGAT GCTGAGAGCTGTTACCTTGTCCACACCCTGCTGGAGTTCTACTTGAAAACTGTTTTCAAAAA CCACCACAATAGAACAGTTGAAGTCAGGACTCTGAAGTCATTCTCTACTCTGGCCAACAACT TTGTTCTCATCGTGTCACAACTGCAACCCAGTCAAGAAAATGAGATGTTTTCCATCAGAGAC AGTGCACACAGGCGGTTTCTGCTATTCCGGAGAGCATTCAAACAGTTGGACGTAGAAGCAGC TCTGACCAAAGCCCTTGGGGAAGTGGACATTCTTCTGACCTGGATGCAGAAATTCTACAAGC TCTGAATGTCTAGACCAGGACCTCCCTCCCCCTGGCACTGGTTTGTTCCCTGTGTCATTTCA AACAGTCTCCCTTCCTATGCTGTTCACTGGACACTTCACGCCCTTGGCCATGGGTCCCATTC TTGGCCCAGGATTATTGTCAAAGAAGTCATTCTTTAAGCAGCGCCAGTGACAGTCAGGGAAG GTGCCTCTGGATGCTGTGAAGAGTCTACAGAGAAGATTCTTGTATTTACTACAACTCTATTT AATTAATGTCAGTATTTCAACTGAAGTTCTATTTATTTGTGAGACTGTAAGTTACATGAAGG CAGCAGAATATTGTGCCCCATGCTTCTTTACCCCTCACAATCCTTGCCACAGTGTGGGGCAG GTTAAAAAACAGAGGGATGCTTGGATGTAAAACTGAACTTCAGAGCATGAAAATCACACT TAAACGATAAAATGTGGATTAAAGTGCCCAGCACAAAGCAGATCCTCAATAAACATTTCATT ${\tt TATCCTAGTCATTCTTCCCTAATCTTCCACTTGAGTGTCAAGCTGACCTTGCTGATGGTGAC}.$ ATTGCACCTGGATGTACTATCCAATCTGTGATGACATTCCCTGCTAATAAAAGACAACATAA CTCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA



></usr/seqdb2/sst/DNA/Dnaseqs.full/ss.DNA88002

><subunit 1 of 1, 206 aa, 1 stop

><MW: 23799, pI: 9.12, NX(S/T): 3

MNFQQRLQSLWTLARPFCPPLLATASQMQMVVLPCLGFTLLLWSQVSGAQGQEFHFGPCQVK GVVPQKLWEAFWAVKDTMQAQDNITSARLLQQEVLQNVSDAESCYLVHTLLEFYLKTVFKNH HNRTVEVRTLKSFSTLANNFVLIVSQLQPSQENEMFSIRDSAHRRFLLFRRAFKQLDVEAAL TKALGEVDILLTWMQKFYKL

Signal sequence:

amino acids 1-42

N-glycosylation sites.

amino acids 85-89, 99-103, 126-130